

The Fate of Liquids Trapped During the Earth's Inner Core Growth

M. Lasbleis, Mathilde Kervazo, G. Choblet

► **To cite this version:**

M. Lasbleis, Mathilde Kervazo, G. Choblet. The Fate of Liquids Trapped During the Earth's Inner Core Growth. *Geophysical Research Letters*, American Geophysical Union, 2020, 47 (2), pp.e2019GL085654. 10.1029/2019GL085654. hal-02481415v2

HAL Id: hal-02481415

<http://hal.univ-nantes.fr/hal-02481415v2>

Submitted on 26 Aug 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Geophysical Research Letters

RESEARCH LETTER

10.1029/2019GL085654

Key Points:

- We calculate the residual melt fraction of liquid in the inner core for different growth scenarios
- An uppermost mushy layer is maintained, of thickness of about 1 km for a viscosity of 10^{20} Pa·s
- Supercooling at the center of the core cannot have been larger than 100 K

Supporting Information:

- Supporting Information S1

Correspondence to:

M. Lasbleis,
marine.lasbleis@univ-nantes.fr

Citation:

Lasbleis, M., Kervazo, M., & Choblet, G. (2020). The fate of liquids

The Fate of Liquids Trapped During the Earth's Inner Core Growth

M. Lasbleis^{1,2}, M. Kervazo¹, and G. Choblet¹

¹Laboratoire de Planétologie et Géodynamique, LPG, UMR 6112, CNRS, Université de Nantes, Université d'Angers, France, ²Earth-Life Science Institute, Tokyo Institute of Technology, Meguro, Japan

Abstract The growth history of the inner core is inherently linked to the thermal history of the Earth. The crystallization of the inner core may have been delayed by supercooling and gone through an initial phase of fast growth after the nucleation barrier has been passed, but so far no evidence or constraint has been proposed to time this possible event. With two-phase flow dynamics, we explore the effect of different growth scenarios for the inner core to determine their effects on its liquid fraction structure. Seismic observations on the melt fraction inside the inner core at present limit the porosity to a maximum of 10% of liquid. Our model provides constraints for the delay in nucleation compared to the time where the first crystal may have started to nucleate, and we find that the supercooling cannot have exceeded 100 K.

Plain Language Summary The magnetic field of the Earth is maintained by the turbulent convection in the liquid outer core. As the inner core grows, latent heat and light elements are released into the liquid outer core, generating upwelling of material and driving the convection. The timing for the nucleation of the inner core, starting the dynamo as we know it today, has been challenging to assess